

UNCLASSIFIED

AD NUMBER
AD868149
NEW LIMITATION CHANGE
TO Approved for public release, distribution unlimited
FROM Distribution authorized to U.S. Gov't. agencies and their contractors; Administrative/Operational Use; APR 1970. Other requests shall be referred to US Army Biological Laboratory, Attn: Technical Release Branch [TID], Fort Detrick, MD 21701.
AUTHORITY
BRL ltr, 29 Sep 1971

THIS PAGE IS UNCLASSIFIED

AD 868149


AD

TECHNICAL MANUSCRIPT 596

**CONCENTRATION OF RIFT VALLEY FEVER VIRUS
BY ALCOHOL PRECIPITATION**

**Frederick Klein
Bill G. Mahlandt
Robert B. Bonner
Ralph E. Lincoln**

APRIL 1970



**DEPARTMENT OF THE ARMY
Fort Detrick
Frederick, Maryland**

Reproduced by the
CLEARINGHOUSE
for Federal Scientific & Technical
Information Springfield Va 22151

49809

OPW	WHITE SECTION	<input type="checkbox"/>
DDO	BUFF SECTION	<input checked="" type="checkbox"/>
UNAN.	GER.	<input type="checkbox"/>
JUSTIFICATION		
BY		
DISTRIBUTION/AVAILABILITY CODES		
DISC.	AVAIL.	NO. OF SPECIAL
2		

Reproduction of this publication in whole or in part is prohibited except with permission of the Commanding Officer, Fort Detrick, ATTN: Technical Releases Branch, Technical Information Division, Fort Detrick, Frederick, Maryland, 21701. However, DDC is authorized to reproduce the publication for United States Government purposes.

DDC AVAILABILITY NOTICES

Qualified requesters may obtain copies of this publication from DDC.

Foreign announcement and dissemination of this publication by DDC is not authorized.

Release or announcement to the public is not authorized.

DISPOSITION INSTRUCTIONS

Destroy this publication when it is no longer needed. Do not return it to the originator.

The findings in this publication are not to be construed as an official Department of the Army position, unless so designated by other authorized documents.

DEPARTMENT OF THE ARMY
Fort Detrick
Frederick, Maryland 21701

TECHNICAL MANUSCRIPT 596

CONCENTRATION OF RIFT VALLEY FEVER VIRUS
BY ALCOHOL PRECIPITATION

Frederick Klein

Bill G. Mahlandt

Hobert E. Bonner

Ralph E. Lincoln

STATEMENT #2 UNCLASSIFIED
This document is subject to special export
controls and each transmittal to foreign
governments or foreign nationals may be made
only with prior approval of Dept. of Army,
Fort Detrick, ATTN: Technical Release Branch/
T10. Frederick, Maryland 21701

Process Development Division
AGENT DEVELOPMENT & ENGINEERING LABORATORIES

Project 1B563603DE32

April 1970

In conducting the research described in this report, the investigators adhered to the "Guide for Laboratory Animal Facilities and Care," as promulgated by the Committee on the Guide for Laboratory Animal Facilities and Care of the Institute of Laboratory Animal Resources, National Academy of Sciences-National Research Council.

ACKNOWLEDGMENT

We thank Jack L. Davis, who was successful in concentrating chikungunya virus by methanol extraction, for his helpful suggestions and encouragement during these studies.

ABSTRACT

Methanol concentrated Rift Valley fever virus approximately 100-fold without appreciable loss of infectivity. The method showed potential both for scale-up and for the concentration of other viruses.

CONCENTRATION OF RIFT VALLEY FEVER VIRUS BY ALCOHOL PRECIPITATION*

Poliovirus preparations have been successfully concentrated by precipitation with methanol and further purified by elution, ultracentrifugation, and enzymatic treatment.^{1,2} They also have been successfully concentrated by chromatography on cellulose ion-exchange columns^{3,4} and on calcium and aluminum phosphate gels.⁵

Earlier, we observed that when Rift Valley fever virus (RVFV) was precipitated with potassium aluminum sulfate (alum),⁶ the volume of viral supernatant fluid was reduced rapidly to 2% of its original volume, yet approximately 100% or more of its original infectivity was retained. In the continuation of these studies, a single-step method for concentration and partial purification of RVFV preparations by precipitation with methanol is described. The method is an adaptation of one that has already proved successful with foot-and-mouth disease virus.⁷ The RVFV particles remained highly infectious after 100-fold concentration.

The origin and maintenance of the wild pantropic van Wyk strain of RVFV⁸ used in this study were described earlier.⁹ Tissue cells (L-DR) grown in suspension to near the peak of the log phase were diluted to approximately 2×10^5 cells/ml in Eagle's minimum essential medium⁹ supplemented with 10% bovine serum. A multiplicity of inoculum of 0.01, seeded directly into the tissue cell cultures, was used. Flasks were routinely incubated at 37 C for 72 hours on a reciprocating shaker (100 3-inch strokes/min).

In concentrating the virus, tissue cells and cellular debris were first removed by low-speed centrifugation. The supernatant fluid was then cooled to -1 C in a dry ice and water mixture, and precooled absolute methanol (-10 C) was added in a ratio of 1:5 (alcohol to virus supernatant). The virus-alcohol mixture was held at -10 C overnight or approximately 16 to 18 hours, after which the mixture was centrifuged for 30 minutes at $6,000 \times g$ in the SS-34 rotor of a Sorvall RC-2 centrifuge operating at -10 C. The supernatant fluid was decanted and the precipitate was resuspended in an equal volume of 199 peptone medium supplemented with 10% bovine serum.

Amount of infective virus was determined by titration in 6- to 8-g Swiss-Webster mice, using four mice per log dilution. The probit method¹⁰ of calculating mouse intracerebral lethal dose₅₀ (MICLD₅₀) values was used. The concentrations achieved by this process were calculated on both per-milliliter and total-volume bases. Results are reported showing both the degree of concentration and percentage recovery of infectivity.

* This report should not be used as a literature citation in material to be published in the open literature. Readers interested in referencing the information contained herein should contact the senior author to ascertain when and where it may appear in citable form.

Results reported here were obtained by using one concentration of alcohol (20%), one holding period (18 hours at -10 C), and one centrifuge speed (2,000 rpm) for the concentration of RVFV particles. This method resulted in concentration of infective virus about 100-fold with little loss of infectivity (Table 1). The supernatant liquid was completely clarified and void of infectious virus particles after one centrifugation, thereby eliminating the time-consuming task of additional centrifugation. Quantitation of the precipitate showed 50 to 80% recovery of infectivity after concentration by this process.

TABLE 1. CONCENTRATION OF RIFT VALLEY FEVER VIRUS
BY ALCOHOL PRECIPITATION

	Original Viral Suspension	Concentrated Viral Suspension	Recovery, per cent
<u>Virus Suspension I</u>			
Repl 1. Volume, ml	100	3.8 ^b /	68
Log ₁₀ MICLD ₅₀ /ml	7.9	9.2	
Log total MICLD ₅₀ ^c	9.9	9.8	
Repl 2. Volume, ml	100	4.2	51
Log ₁₀ MICLD ₅₀ /ml	7.9	9.0	
Log total MICLD ₅₀	9.9	9.6	
<u>Virus Suspension II</u>			
Repl 1. Volume, ml	100	2.8	73
Log ₁₀ MICLD ₅₀ /ml	7.8	9.2	
Log total MICLD ₅₀	9.8	9.6	
Repl 2. Volume, ml	100	3.1	81
Log ₁₀ MICLD ₅₀ /ml	7.8	9.2	
Log total MICLD ₅₀	9.8	9.7	
Repl 3. Volume, ml	100	3.5	62
Log ₁₀ MICLD ₅₀ /ml	7.8	9.0	
Log total MICLD ₅₀	9.8	9.6	

- a. For all five cultures, supernatant volume was 100 ml. Likewise, value for all five supernatants was $<10^{-1}$ MICLD₅₀/ml, total MICLD₅₀ and concentration.
- b. Volume represents the original precipitate plus an equal volume of suspending medium.
- c. Volume x MICLD₅₀/ml = total MICLD₅₀.

This concentration method has the advantages of simplicity and potential scale-up for any volume of culture, and our results on concentration achieved and percentage loss of infectivity seem very promising. Certain possibilities for its use are: (i) the reduction of large volumes of virus suspensions without appreciable loss of infectivity, thereby making storage and further purification possible; (ii) the preparation of antigens (live, dead, or attenuated) for vaccine production; and (iii) if some purification is achieved, the removal of potentially harmful and/or interfering substances originating from destruction of the tissue cells. We are presently attempting to consider these possibilities and to assess the effect of continuous centrifuge flow instead of batch treatment utilizing this concentration process. This method also seems to offer potential for concentrating other viruses that contain essential lipoid material and are sensitive to ether or other fat solvents.

LITERATURE CITED

1. Mayer, M.M.; Rapp, H.J.; Roizman, B.; Klein, S.W.; Cowan, K.M.; Lukens, D.; Schwerdt, C.E.; Schaffer, F.L.; Charney, J. 1957. The purification of poliomyelitis virus as studied by complement fixation. *J. Immunol.* 78:435-455.
2. Schwerdt, C.E.; Schaffer, F.L. 1956. Purification of poliomyelitis virus propagated in tissue culture. *Virology* 2:665-678.
3. Hayer, B.H.; Bolton, E.T.; Ritter, D.B.; Ribí, E. 1959. Simple method for preparation of purified radioactive poliovirus particles: Electron micrograph. *Virology* 7:462-470.
4. Taylor, J.; Graham, A.F. 1958. Purification of poliovirus labelled with radiophosphorus. *Virology* 6:498-498.
5. Taverne, J.; Marshall, I.H.; Fulton, F. 1957. The purification and concentration of viruses and virus soluble antigens on calcium phosphate. *J. Gen. Microbiol.* 19:451-461.
6. Klein, F.; Mahlandt, B.G.; Cockey, R.R.; Lincoln, R.E. 1970. Concentration of Rift Valley fever and chikungunya viruses by precipitation with ammonium sulfate and potassium aluminum sulfate, (Technical Manuscript 598). Process Development Division, Fort Detrick, Frederick, Maryland.
7. Bachrach, H.L.; Trautman, R.; Breese, S.S., Jr. 1964. Chemical and physical properties of virtually pure foot-and-mouth disease virus. *Amer. J. Vet. Res.* 25:333-342.
8. Kaschula, V.R. 1953. The propagation and modification of strains of Rift Valley fever viruses in embryonated eggs and their use as immunizing agents for domestic ruminants. Thesis, Doctor of Veterinary Science, University of Pretoria, South Africa.
9. Klein, F.; Walker, J.S.; Mahlandt, B.G.; Carter, R.C.; Orlando, M.D.; Weirether, F.J.; Lincoln, R.E. 1969. Interacting factors that influence long-term storage of live Pasteurella tularensis vaccine and Rift Valley fever virus. *Appl. Microbiol.* 17:427-434.
10. Finney, D.J. 1952. Statistical methods in biological assay, p. 524-530. Hafner Publishing Company, New York.

Unclassified

9

Security Classification

DOCUMENT CONTROL DATA - R & D		
<small>(Security classification of title, body of abstract and indexing annotation must be entered when the overall report is classified)</small>		
1. ORIGINATING ACTIVITY (Corporate author)		2a. REPORT SECURITY CLASSIFICATION
Department of the Army Fort Detrick, Frederick, Maryland, 21701		Unclassified
		2b. GROUP
3. REPORT TITLE		
CONCENTRATION OF RIFT VALLEY FEVER VIRUS BY ALCOHOL PRECIPITATION		
4. DESCRIPTIVE NOTES (Type of report and inclusive dates)		
5. AUTHOR(S) (First name, middle initial, last name)		
Frederick (NMI) Klein Hobert B. Bonner Bill G. Mahlandt Ralph E. Lincoln		
6. REPORT DATE	7a. TOTAL NO. OF PAGES	7b. NO. OF REFS
April 1970	9	10
8a. CONTRACT OR GRANT NO.		8b. ORIGINATOR'S REPORT NUMBER(S)
a. PROJECT NO 1B563603DE32		Technical Manuscript 596
c. Task-Work Unit 04-003		9b. OTHER REPORT NUM(S) (Any other numbers that may be assigned this report)
d. DD 1498 Agency Access. DA OL 0305		CMs 6663 AMXFD-AE-T 49809
10. DISTRIBUTION STATEMENT		
Qualified requesters may obtain copies of this publication from DDC. Foreign announcement and dissemination of this publication by DDC is not authorized. Release or announcement to the public is not authorized.		
11. SUPPLEMENTARY NOTES		12. SPONSORING MILITARY ACTIVITY
Process Development Division		Department of the Army Fort Detrick, Frederick, Md., 21701
13. ABSTRACT		
Methanol concentrated Rift Valley fever virus approximately 100-fold without appreciable loss of infectivity. The method showed potential both for scale-up and for the concentration of other viruses.		
14. Key Words		
Rift Valley fever virus Methanol Concentration		

DD FORM 1473

REPLACES DD FORM 1473, 1 JAN 64, WHICH IS OBSOLETE FOR ARMY USE.

Unclassified
Security Classification